



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

which was later identified for me by Mr. Frank M. Chapman, to whom I sent it, as *Melospiza lincolni*.

This, I believe, is the first known instance of the occurrence of the species in New Brunswick.—WILLIE H. MOORE, *Scotch Lake, N. B.*

Rank of the Sage Sparrow.—On page 58 of the current volume of 'The Auk,' Mr. Joseph Grinnell states that he found *Amphispiza belli* and *Amphispiza belli nevadensis* inhabiting the same locality at the head of the Little Tujunga Cañon, Los Angeles County, California, in July, 1897. This area is on the western slope of the divide, though not more than ten miles from the Mohave Desert. Mr. Grinnell further states that he has never learned of any intermediate specimens between the two forms, and consequently argues that they are specifically distinct.

On the Death Valley Expedition in 1891, Mr. Frank Stephens collected a number of specimens on the eastern slope of the Sierras, opposite the south end of Owens Lake, which I reported as being intermediate in color and size (N. Am. Fauna, No. 7, p. 98).

Taking this into consideration and the fact that *Amphispiza b. nevadensis* had evidently wandered from their desert home, as Mr. Grinnell writes me they were fully fledged, I cannot agree with him that there is any reason for considering the two forms more than subspecifically distinct.—A. K. FISHER, *Washington, D. C.*

Wintering of the Towhee (*Pipilo erythrophthalmus*) at Rockaway Beach, L. I.—On the 29th of January, 1898, I made a collecting trip to Rockaway Beach in the hope of seeing some winter birds, as the weather had been very cold for several days, and on the day in question the air was full of flying snow. While passing through a small thicket of brambles I felt sure I heard the Towhee's note, and started in to investigate. Although he was exceedingly wild, I at length caught a glimpse of him, and by remaining quiet for some time eventually secured him. Later in the day, I found three more, all males, as was the one I shot. It is quite evident that this bird occasionally winters much further north than is generally supposed, as there are also records from Longwood, Mass., on Christmas, and Bedford, Mass., on Jan. 2 (Auk, July, 1896).

Mr. L. S. Foster informs me that he secured a specimen on Feb. 22, near Oradell, N. J.—HARRY WEBB FLOYD, *New York City.*

The Rose-breasted Grosbeak in California.—During a collecting trip last summer in northern California, Dr. C. H. Gilbert and a party of students secured some birds that were new to the fauna of the State. While at Meyer's, Humboldt Co., Cal., July 1, 1897, the attention of the party was attracted by a string of strange birds that had been shot sometime before and were already in the early stages of decomposition. Not being able to decide what the birds were, several heads were cut off,